

**End-line Survey on Scaling up the Use of Zinc and  
Oral Rehydration Salts (ORS) in the Treatment of Diarrhoea  
among the 6-59 Months Children to Reduce Child  
Morbidity and Mortality in Bangladesh**

**(Contract no: 10-1531-HUMDEV-26)**

*Prepared for*

**Nutrition International**

Commercial Cove,  
House: 56/C (Level 8)  
Road 132, Gulshan – 1,  
Dhaka 1212, Bangladesh

*Prepared by*



**Human Development Research Centre**

House 5, Road 8, Mohammadia Housing Society,  
Mohammadpur, Dhaka 1207, Bangladesh  
Phone: (88 02) 58150381, 58157621, 8101704, Fax: (88 02) 58157620  
E-mail: [info@harc-bd.com](mailto:info@harc-bd.com), Web: [www.hdrc-bd.com](http://www.hdrc-bd.com)

August, 2017

## Acknowledgments

Micronutrient Initiative (MI), Bangladesh demonstrated a model with promising result to improve child survival through strengthening zinc supplementation and ORS distribution in the treatment of diarrhoea in children 6-59 months. Currently, MI is scaling up the best practices of the demonstration program in 9 districts in request of Primary Health Care under Directorate General of Health Services. Three out of these first year nine scale up project districts are being evaluated using a pre and post intervention survey design with comparison districts. A baseline survey of the program has been undertaken in early 2015 in 3 intervention districts (Kishorejganj, Bogra and Patuakhali) and three comparison districts (Brahmanbaria, Noagaon and Pirojpur). Now, MI has commissioned an 'End-line Survey on Scaling up the Use of Zinc and Oral Rehydration Salts (ORS) in the Treatment of Diarrhoea among the 6-59 Months Children to Reduce Child Morbidity and Mortality in Bangladesh' in those districts which has been undertaken by Human Development Research Centre (HDRC).

We are grateful to the relevant officials of MI (currently Nutrition International) for entrusting us to carry out this important end-line survey of the project.

We are grateful to Dr. Mustafizur Rahman, Former Country Director, MI Country Office in Bangladesh, and Mr. Zaki Hasan, current Country Director, Nutrition International (former MI) for their enthusiasm towards the study. We are also grateful to Dr. Khairul Islam, National Program Officer, and Ms. Samsun Nahar, Nutrition International (former MI), Bangladesh for their stimulating inputs, and unstinted support at different stages of this survey.

We are indebted to Monoj Kumar Raut, Ph.D., Regional Manager, Research and Evaluation, Nutrition International, Asia from the Regional Office, India, and Mr. Kirti Warvadekar, from Head office of Nutrition International for their invaluable research inputs and monitoring support towards the study at different stages.

We are truly indebted to Daniel Lopez De Romana, Ph.D., Technical Advisor, Infant and Young Child Health and Nutrition for his valuable comments made on the draft report.

We are grateful to all the caregiver/mothers of the diarrhoea children, Health Workers, Health supervisors, *Gram Dakters* and Health Managers who provided us a lot of valuable information on the subject.

Above all, we will remain ever grateful to the in-house staff members of HDRC who worked untiringly at all the stages, specially, Ms. Nurunnahar for her valuable contribution. We thank all those lovely and uncomplaining souls at HDRC. The successful administration of this end-line survey would not have been possible without the commitment and dedication of all those who were involved in this process.

Dr. Md. Murtaza Majid  
Principal Investigator and Team Leader  
Human Development Research Centre (HDRC)

Dhaka: August 2017

## ABBREVIATIONS

AHI	Assistant Health Inspector
BCI	Behaviour Change Intervention
CC	Community Clinic
CHCP	Community Health Care Provider
CS	Civil Surgeon
DCI	Data Collection Instrument
DDFP	Deputy Director of Family Planning
DGHS	Directorate General of Health Services
FPI	Family Planning Inspector
FWA	Family Welfare Assistant
FWV	Family Welfare Visitor
GoB	Government of Bangladesh
HA	Health Assistant
HH	Household
HI	Health Inspector
HSC	Higher Secondary School Certificate
IMCI	Integrated Management of Childhood Illness
IPC	Inter Personal Communication
IRB	International Research Bureau
KAP	Knowledge, Attitude and Practice
MI	Micronutrient Initiative
ORS	Oral Rehydration Salt
PHC	Primary Health Care
SACMO	Sub-assistant Community Medical Officer
SMS	Short Message Service
SSC	Secondary School Certificate
UFPO	Upazila Family Planning Officer
UH&FPO	Upazila Health and Family Planning Office
UHC	Upazila Health Complex
UNICEF	United Nations Children's Fund
WHO	World Health Organization

# CONTENTS

Sl. No.	Title	Page #
	<b>Executive Summary</b> .....	1-v
	<b>CHAPTER-1: INTRODUCTION</b> .....	<b>1</b>
1.1	Background .....	1
1.2	Objectives of the End-line Survey .....	2
1.3	Key research questions .....	2
1.4	Organization of the report.....	3
	<b>CHAPTER-2: STUDY DESIGN AND APPROACH</b> .....	<b>4</b>
2.1	Methodology .....	4
2.2	Study respondents and Data collection instruments (DCIs) .....	4
2.3	Study locations .....	4
2.4	Sample Size & Sample Selection Criteria.....	5
2.4.1	Sample size for Care-givers of child with diarrhoea in the previous month.....	5
2.4.2	Study among Health managers, Health supervisors, Health workers and <i>Gram Dakters</i> .....	5
2.5	Data Analysis .....	6
	<b>CHAPTER 3: SOCIO-DEMOGRAPHIC CHARACTERITICS</b> .....	<b>8</b>
3.1	Household Characteristics .....	8
3.1.1	Age of the household members.....	8
3.1.2	Sex ratio .....	9
3.1.3	Marital status.....	9
3.1.4	Education .....	9
3.1.5	Source of Household Income.....	9
3.1.6	Average Monthly Household Income .....	9
3.1.7	Occupation of main wage earner .....	10
3.1.8	Main source of drinking water .....	10
3.1.9	Type of sanitation facility .....	10
3.1.10	Use of fuel for cooking .....	10
3.1.11	Functional household assets.....	10
3.1.12	Housing Characteristics .....	11
3.1.13	Power supply.....	11
3.1.14	Membership with credit group.....	12
3.2	Respondents characteristics .....	12
3.2.1	Relationship of with the HH head.....	12
3.2.2	Religion.....	12
3.2.3	Education of respondents.....	13
3.2.4	Education of husband.....	13
3.2.5	Exposure to media .....	13
	<b>CHAPTER 4: DIARRHOEAL EPISODE AND UNDERSTANDING OF THE MOTHERS/ CARETAKERS ON DIARRHOEA</b> .....	<b>15</b>
4.1	Age and sex of the Children.....	15
4.2	Episodes of Childhood Diarrhoea.....	16
4.3	Knowledge of the mother/caregiver about diarrhoea.....	17
4.4	Knowledge on key manifestation on dehydration.....	17
	<b>CHAPTER 5: KNOWLEDGE AND ATTITUDE OF MOTHERS/CAREGIVERS TOWARDS ORS AND ZINC</b> .....	<b>19</b>
5.1	Knowledge about ORS .....	19
5.2	Knowledge about the source of ORS.....	19
5.3	Knowledge about Zinc Tablet.....	20

Sl. No.	Title	Page #
5.4	Knowledge about Sources of Zinc Tablet.....	20
5.5	Knowledge about Duration of Treatment with Zinc Tablet.....	21
5.6	Knowledge about Benefit of Zinc Supplement.....	22
<b>CHAPTER 6: DIARRHOEAL EPISODE AND UNDERSTANDING OF THE MOTHERS/ CARETAKERS ON DIARRHOEA .....</b>		<b>23</b>
6.1	Time of Seeking Treatment .....	23
6.2	Where and Whom Consulted for Treatment.....	23
6.3	Prescribed Treatment for Acute Childhood Diarrhoea .....	25
6.4	Time Taken to Control Diarrhoea.....	25
6.5	Time Taken to Complete Cure from Diarrhoea Related Illness .....	26
6.6	Compliance of Mothers/Caregivers to Prescribed Treatment.....	26
6.7	Supply and consumption of ORS.....	28
6.8	Place of Collection of ORS.....	29
6.9	Practice of using of Zinc Supplementation .....	30
6.10	Recommended course and compliance to zinc tablet .....	31
6.11	Compliance/Adherence to Zinc Syrup .....	34
6.12	Place of Collection of Zinc Tablet and/or Syrup .....	35
6.13	Acceptability of Zinc Tablet and Syrup.....	37
6.14	Side Effects of Zinc and Management of Side Effects .....	38
6.15	Future Intention of Using Zinc .....	39
6.16	Encouraging Others for Use of Zinc .....	39
6.17	Home Visit by the Health Workers.....	40
6.18	Counseling by the Health Workers .....	40
6.19	Contact with the Health Worker .....	40
<b>CHAPTER-7: KNOWLEDGE, PERCEPTION AND ACTIVITIES OF THE HEALTH WORKERS.....</b>		<b>42</b>
7.1	Background Characteristics of the Health Workers.....	42
7.1.1	Type of the Health Workers.....	42
7.1.2	Age and Sex of the Health Workers .....	42
7.1.3	Education of the Health Workers.....	42
7.1.4	Receipt of Basic Training as Health Workers .....	43
7.1.5	Receipt of Training on Use of Zinc in Diarrhoea from IMCI programme of DGHS or MI .....	43
7.1.6	Frequency and Timing of Receipt of Training on Zinc in Diarrhoea .....	44
7.1.7	Receipt of Refresher Training.....	44
7.2	Catchment Population and Client Served by the Health Workers .....	44
7.2.1	Knowledge on Diarrhoea .....	44
7.2.2	Knowledge on Manifestations of Dehydration .....	45
7.2.3	Common Place for Consultation in Childhood Diarrhoea .....	46
7.2.4	Knowledge on Benefits of Zinc Supplement in Acute Severe Diarrhoea.....	47
7.3.	Knowledge of Continuation of Zinc Supplement for 10 days and reasons thereof .....	47
7.3.1	Knowledge on Dosage of Zinc .....	48
7.3.2	Knowledge on Liquids Used to Dissolve Zinc Tablet before Administration.....	49
7.3.3	Knowledge and Demonstration of Preparation of ORS.....	50
7.3.4	Demonstration of Procedure of Administration of Dispersible Zinc Tablet.....	50
7.4	Opinion Regarding Efficacy of Zinc with ORS and Future Intention of Using Zinc .....	51
7.5	Practice of Health Workers in the Management of Childhood Diarrhoea .....	51
7.5.1	Practice of Zinc Supplement in Last One Month.....	52
7.5.2	Instructions on Zinc Supplement .....	53
7.6	Household Visit Made by the Health Workers .....	53
7.7	Status of Stock Register and Reporting Form.....	54
7.8	Receipt and Distribution of Zinc Blister Packs at the Health Centre.....	55

Sl. No.	Title	Page #
7.9	Receipt and Distribution of ORS Sachets at the Health Centre .....	55
7.10	Childhood Diarrhoea seen at Last Month .....	56
7.11	Status of Anti-diarrhoeal Management in Last Month .....	56
7.12	Stock-out and Stock-in at Present of Zinc and ORS .....	57
7.13	Knowledge on How to Estimate Demand of Zinc and ORS in Respective Area .....	58
7.14	Problems in Supply and Means of Way Out.....	59
7.15	Receipt of Communication/IPC Material under this Project .....	59
7.16	Conduction of Group Counseling Session .....	60
7.17	Counseling by the Health Workers: Reporting made by Mothers/Care-givers .....	60
<b>CHAPTER 8: KNOWLEDGE, PERCEPTION AND ACTIVITIES OF HEALTH SUPERVISORS .....</b>		<b>61</b>
8.1	Demographics of Health Supervisors .....	61
8.1.1	Types of the Health Supervisors Interviewed.....	61
8.1.2	Age and Sex of the Health Supervisors .....	61
8.1.3	Education of the Health Supervisors.....	62
8.2	Duties and Responsibilities of the Supervisors.....	62
8.3	Knowledge and Skills of the Supervisors Related to Diarrhoea Management .....	63
8.3.1	Expected Number of Cases of Diarrhoea in Catchment area in each Month .....	63
8.3.2	Knowledge about the Program on Use of Zinc and ORS in Management of Diarrhoea ..	63
8.3.3	Knowledge on the Correct Dosage of Zinc Tablets.....	63
8.3.4	Knowledge of Correct Duration of Zinc Tablets Intake .....	64
8.3.5	Knowledge on the Benefits of Giving Zinc in Acute Diarrhoea.....	64
8.3.6	Knowledge on the Reason for Intake of Zinc Tablet for a period of 10 Days even if Diarrhoea Stops.....	65
8.4	ORS and Zinc Supplies and Stock.....	65
8.4.1	Stock out for Zinc and ORS in their Catchment Area in the Last Month.....	65
8.4.2	Steps to Overcome the Stock out Situation.....	65
8.5	Monitoring and Supervision .....	65
8.5.1	Frequency of Visits to Undertake Supervision/Monitoring.....	65
8.5.2	Using tools during Supervising/monitoring visits .....	66
8.5.3	Use of Checklist/Monitoring Tools during Monitoring Visits.....	66
8.5.4	Motivating Staff to Seek out Diarrhoea Cases through Increased Home Visits .....	66
8.5.5	Conducting Home Visit .....	67
8.5.6	Monitoring during Home Visits.....	67
8.5.7	Use of BCC Materials during Home Visits or Supervision Staff .....	67
8.5.8	Types of BCC/IEC Materials Used During Home Visit or Supervision of Staffs.....	67
8.5.9	Counseling the Caregivers to comply with the full Course and Benefits of Zinc.....	68
8.5.10	Monthly Report Compilation by Supervisors .....	68
8.5.11	Regularity in Receiving HMIS/Monitoring Reports on Zinc and ORS.....	68
8.5.12	Sharing Feedback of Monitoring Visit to Staffs .....	68
8.5.13	Sharing Key Observations from Monitoring Visits in the Monthly Meeting .....	69
8.6	Training Related to ORS and Zinc.....	69
8.7	Suggestion for Improved Coverage and Compliance of Zinc.....	69
<b>CHAPTER- 9: KNOWLEDGE, PERCEPTION AND ACTIVITIES OF THE GRAM DAKTARS. 70</b>		
9.1	Knowledge and Skills of the <i>Gram Daktars</i> Related to Diarrhoeal Management.....	70
9.2	Training of <i>Gram Daktars</i> on ORS and Zinc .....	71
9.3	Pattern of Treatment for Childhood Diarrhoea in the Last Week .....	71
9.4	Receipt of Inter Personal Communication (IPC) tool(s) from program .....	72
9.5	Availability of zinc tablets in pharmacy in the locality/neighborhood.....	73
9.6	Suggestions to improve coverage and compliance of zinc and ORS.....	73

Sl. No.	Title	Page #
<b>CHAPTER- 10: OPINION AND SUGGESTIONS OF THE HEALTH MANAGERS .....</b>		<b>74</b>
10.1	Status of Zinc supplementation .....	74
10.2	Training of the frontline workers .....	74
10.3	Role of counseling by the frontline workers .....	75
10.4	Opinion about the role of ensuring adequate supplies .....	75
10.5	Monitoring or supervision of program activities including stock of Zinc and ORS.....	75
10.6	Discussion issues about the program in monthly coordination meeting.....	76
10.7	Suggestions on the IEC/ BCC programme .....	76
10.8	Suggestions on programme implementation to improve the coverage and compliance of Zinc and ORS among diarrheal children.....	77
Reference	.....	78

### *List of Figures*

Figure 3.1:	Percentage distribution of age of all members of the households.....	8
Figure 3.2:	Percentage distribution of respondents regarding main source of income of the households .....	9
Figure 3.3:	Percentage distribution of respondents regarding functional assists under the ownership of the household at present.....	10
Figure 3.4:	Percentage distribution of respondents regarding source of electricity in the households .....	12
Figure 3.5:	Percentage distribution of the household members regarding microcredit organization in which having membership .....	12
Figure 3.6:	Percentage distribution of respondents regarding which sites usually go to .....	14
Figure 4.1:	Distribution of under-five children by age in months suffered from acute diarrhoea 30 days prior tp survey in baseline and End-line survey (%) .....	15
Figure 4.2:	Distribution children aged 6-59 months by number of diarrhoeal episode in the last 30 days prior to the survey in intervention and comparison areas (%) .....	16
Figure 4.3:	Distribution of mothers/caregivers by their knowledge on diarrhoea by frequency of loose stool in intervention and comparison areas (%) .....	17
Figure 5.1:	Distribution of mothers/caregivers by their correct knowledge about preparation ORS in intervention and comparison areas at baseline and End-line survey and the change between the periods (%) .....	19
Figure 5.2:	Distribution of mothers/care givers by their knowledge on zinc tablet in intervention and comparison areas at baseline and End-line (%) .....	20
Figure 5.3:	Distribution of mothers/care givers by their knowledge on duration of zinc tablet supplementation in intervention and comparison areas at baseline and End-line (%).....	22
Figure 6.1:	Distribution of mothers/caregivers by administration of zinc tablet and syrup .....	30
Figure 6.2:	Distribution of the children by acceptability of zinc tablet and syrup (%) .....	38
Figure 6.3:	Distribution of mothers/caregivers by intention to future use of zinc (%) .....	39

### *List of Tables*

Table 3.1:	Percentage distribution of household by number of rooms for dwelling purpose .....	11
Table 3.2:	Percentage distribution of respondents regarding her religion .....	13
Table 3.3:	Percentage distribution of respondents regarding highest class passed.....	13
Table 4.1:	Percentage distribution of children by age (in month) who suffered from diarrhoea in last 30 days prior to the survey .....	15
Table 4.2:	Percentage distribution of the children aged 6-59 months by the duration of diarrhoea in the last episode within last 30 days prior to the survey .....	16
Table 4.3:	Percentage distribution of the mothers/caregivers by their knowledge about signs and symptoms of dehydration at baseline and end-line including change of knowledge.....	18

Sl. No.	Title	Page #
Table 5.1:	Percentage distribution of mothers/caregivers by knowledge about sources of ORS across the public, private and other sectors at baseline and end-line.....	20
Table 5.2:	Percentage distribution of mothers/caregivers of under-five children by knowledge on source of zinc tablet across the sectors at baseline and end-line .....	21
Table 5.3:	Percentage distribution of mothers/caregivers by knowledge on benefit of zinc supplement in acute diarrhea at baseline and end-line.....	22
Table 6.1:	Percentage distribution of the mothers/caregivers by time of seeking treatment at the onset of acute diarrhoea of their children .....	23
Table 6.2:	Percentage distribution of the mothers/caregivers of children, who have sought care for diarrhoea from any source, public source, and private source by baseline and end-line.....	24
Table 6.3:	Percentage distribution children who had diarrhoea in last 30 days by type of treatment prescribed by the service provider at baseline and end-line .....	25
Table 6.4:	Percentage distribution children with acute diarrhoea by time taken to recover from diarrhoea .....	26
Table 6.5:	Percentage distribution of the children by the duration of diarrhea in the last episode of diarrhea before 30 days of survey.....	26
Table 6.6:	Percentage distribution of the mothers/caregivers by type of treatment administered to the children during the last episode of diarrhoea within 30 days .....	27
Table 6.7:	Percentage distribution of ORS sachets by number of sachets distributed to and duration of consumption of ORS per child at the time of diarrhoea .....	28
Table 6.8:	Percentage distribution mothers/caregivers by major sources of ORS collection .....	29
Table 6.9:	Association of socio-demographic variables with compliance to zinc course of therapy .....	32
Table 6.10:	Percentage distribution of the mothers/caregivers by number of zinc tablet given to their children per day during last diarrhea 30 days prior to survey.....	33
Table 6.11:	Percentage distribution of the mothers/caregivers by their common practice of dissolving the zinc tablet with water before administering to child .....	33
Table 6.12:	Percentage distribution of the mothers/caregivers by number of teaspoonful of zinc syrup given to their children in a day .....	35
Table 6.13:	Percentage distribution of the mothers/caregivers by main places of collection of zinc tablet .....	36
Table 6.14:	Percentage distribution of the mothers/caregivers by main places of collection of zinc syrup .....	36
Table 6.15:	The estimated coefficient and odds ratio of the binary regression for coverage and adherence of zinc .....	37
Table 6.16:	Percentage distribution of the mothers/caregivers by practice of administering zinc tablet if disliked by the children .....	38
Table 6.17:	Percentage distribution of the children by types of side effects experienced .....	39
Table 6.18:	Percentage distribution of the health workers who visited mothers/caregivers house for counseling on diarrhoeal management even their children were not suffering with diarrhoea .....	40
Table 6.19:	Percentage distribution of the mothers/caregivers counseled by the health workers by key subject matters related to diarrhoea .....	40
Table 6.20:	Percentage distribution of the mothers/caregivers who have the contact address/number of the health workers for future communication .....	41
Table 7.1:	Percentage distribution of health workers by sex .....	42
Table 7.2:	Percentage distribution of health workers by their educational qualification.....	43
Table 7.3:	Percentage distribution of health workers by how many months back they have attended in basic training.....	43
Table 7.4:	Percentage distribution of health workers by receipt of training from IMCI, DGHS or MI on use of zinc.....	43
Table 7.5:	Percentage distribution of health workers by their knowledge about diarrhoea.....	45

Sl. No.	Title	Page #
Table 7.6:	Percentage distribution of health workers by their perception about diarrhea in terms of frequency of loose motion per day .....	45
Table 7.7:	Percentage distribution of health workers by their knowledge on common manifestations of dehydration.....	46
Table 7.8:	Percentage distribution of health workers by their opinion regarding the common health seeking place of the community people for the treatment childhood diarrhea .....	46
Table 7.9:	Percentage distribution of health workers by their knowledge on benefits of dispersible zinc tablet .....	47
Table 7.10:	Percentage distribution of health workers by their knowledge about duration of continuation zinc tablet without interruption.....	48
Table 7.11:	Percentage distribution of health workers by their knowledge on recommended dosage of Zinc tablets according to the age of the children.....	49
Table 7.12:	Percentage distribution of health workers by their knowledge about type of liquid is usually mixed with zinc tablet before giving to the children.....	50
Table 7.13:	Percentage distribution of health workers whether they able to demonstrate the method of using of dispersible zinc tablet correctly .....	51
Table 7.14:	Percentage distribution of health workers by type of treatment given the children during the last diarrhoeal episode .....	52
Table 7.15:	Percentage distribution of health workers by whether zinc tablet is given to anyone in the past one month.....	52
Table 7.16:	Percentage distribution of health workers by whether they advice to start zinc and ORS as a first line treatment of childhood diarrhoea .....	53
Table 7.17:	Percentage distribution of health workers by home visit of diarrhoea affected children.....	53
Table 7.18:	Percentage distribution of health workers by their information on availability stock registers and reporting forms at the health facilities.....	55
Table 7.19:	Average number of zinc blister packs received by the health workers in the last month.....	55
Table 7.20:	Average number of zinc blister packs distributed by the health workers in the last month ..	55
Table 7.21:	Average number of ORS sachets received by the health workers in the last month .....	56
Table 7.22:	Average number of ORS sachets distributed by the health workers in the last month.....	56
Table 7.23:	Number of children with diarrhoea seen by the health workers in last month .....	56
Table 7.24:	Percentage distribution of health workers who could not give zinc tablet to the client due to ran out of supply in the last month .....	57
Table 7.25:	Current stock of zinc blister pack (in number) of the health worker .....	58
Table 7.26:	Current stock of ORS sachets (in number) of the health worker .....	58
Table 7.27:	Percentage distribution of health workers by adequate stock of ORS and zinc at present ....	58
Table 7.28:	Percentage distribution of health workers by knowledge about how to calculate demand estimation of zinc and ORS .....	58
Table 7.29:	Percentage distribution of health workers by experience of problem and way to resolve ....	59
Table 7.30:	Percentage distribution of the health workers by conduction of group counseling sessions in the last month.....	60
Table 7.31:	Percentage distribution of the health workers whether discuss the benefit of zinc in group counseling session in the last month.....	60
Table 8.1:	Percentage distribution of health supervisors interviewed by their age.....	61
Table 8.2:	Percentage distribution of health workers by sex .....	62
Table 8.3:	Percentage distribution of supervisors by their Education .....	62
Table 8.4:	Percentage distribution of supervisors by their job responsibilities .....	63
Table 8.5:	Percentage distribution of supervisors by their knowledge about the program on use of zinc and ORS in management of diarrhoea .....	63

Sl. No.	Title	Page #
Table 8.6:	Percentage distribution of supervisors by their knowledge on the correct dosage of zinc tablets .....	64
Table 8.7:	Percentage distribution of supervisors by their knowledge of correct duration of Zinc tablets intake .....	64
Table 8.8:	Percentage distribution of supervisors by their knowledge on the benefits of giving Zinc in acute diarrhoea.....	64
Table 8.9:	Percentage distribution of supervisors by frequency of visits to undertake supervision / monitoring .....	66
Table 8.10:	Percentage distribution of supervisors by whether they use any checklist/monitoring tool during monitoring visit .....	66
Table 8.11:	Percentage distribution of supervisors by whether they motivate staff to seek out diarrhoea cases through increased home visits .....	67
Table 8.12:	Percentage distribution of supervisors by whether they use any BCC materials during home visit or during the supervision of their staff.....	67
Table 8.13:	Percentage distribution of supervisors by whether they regularly receive HMIS/ monitoring reports on zinc and ORS .....	68
Table 8.14:	Percentage distribution of supervisors by whether they need any further training.....	69
Table 9.1:	Knowledge and skills of Gram Daktars in relation to diarrhoea management .....	71
Table 9.2:	Treatment behaviors of <i>Gram Daktars</i> for childhood diarrhoea in past one week.....	72
Table 9.3:	Percentage distribution of <i>Gram Daktars</i> by treatment with zinc and ORS for childhood diarrhoea .....	72
Table 9.4:	Percentage distribution of Gram Daktars by whether zinc tablets are available in pharmacy in the locality/neighborhood area.....	73

## ANNEXUE

Annex-1:	Data Tables .....	80-231
Annex-2:	Data Collection Instruments (DCIs) .....	232-269
Annex-3:	Study Team Members.....	270-271

# Executive Summary

## Background

In the context of high Infant and Under Five Mortality Rate, Micronutrient Initiative (MI), Bangladesh demonstrated a model for strengthening zinc supplementation and ORS distribution in the treatment of diarrhoea in children 6-59 months to improve child survival. In view of the promising results of the demonstration program, MI is scaling up the best practices of the demonstration program in additional districts in request of Primary Health Care under Directorate General of Health Services (PHC, DGHS). The scale up program will be implemented in 27 districts in a phased manner with nine districts being initiated in the first year of the program. Three out of these first year nine scale up project districts are being evaluated using a pre and post intervention survey design with comparison districts to allow for maximum program exposure. A baseline survey of the programme has been undertaken in early 2015 in three intervention districts (Kishoreganj, Bogra and Patuakhali) and three comparison districts (Brahmanbaria, Noagaon and Pirojpur). Now, an end-line survey with key objectives to measure the knowledge, attitude and practices (KAP) regarding childhood diarrhea management in those three intervention and three comparison districts has been commissioned by MI (currently Nutrition International) and been undertaken by Human Development Research Centre (HDRC).

*The specific objectives of the end-line survey are to:*

1. Estimate coverage and compliance of Zinc supplementation with ORS during diarrhoea episodes of children 6-59 months of age
2. Assess the knowledge, attitude and practices among mothers and health workers in treatment of childhood diarrhoea; and
3. Assess the knowledge, skill and behavior of health service providers related to services and advice provided to caregivers of 6-59 months old children suffering from diarrhoea.

## Methodology

The current study is for end-line estimates of a quasi-experimental pre and post-intervention repeated cross sectional survey design for collecting quantitative data. The study was conducted in three purposively selected MI Intervention districts together with three comparison non-project districts adjacent to the selected program districts having similar socio-economic and health characteristics. In order to compare findings from baseline and end line data, the sample size, sample locations (District, Upazila, Union) and sample selection strategy during end line was identical to the baseline.

Tabulation for quantitative data analysis techniques mostly includes uni-variate analysis and bi-variate analysis. The quantitative analysis was summarized by districts and intervention-Control groups. Difference-in-difference (DiD) technique was applied to assess the net impact of project intervention. Also statistical tests (hypothesis testing) were performed to assess significance of the changes. Qualitative interviews were scripted and then content analysis was performed. Necessary triangulations were also performed.

## Study Findings

### Background situation of mothers/caregivers

**Household characteristics:** Around one-fourth household members are below 5 years age, with mean age of the under-five children 30 months. More than two-fifth of the household members are under 18 years. Among the household members, almost half are married, and of the rest, 47 percent are unmarried, and 2 percent widow/widower. The average monthly household income is Taka 12,475 in intervention and Taka 13,143 in comparison area. Main sources of household income are- day laborer, followed by agriculture and business. Main source of drinking water is tube-well in more than 90 percent households.

**Education, Exposure to media, and Membership with credit group:** More than 85 percent mothers/caregivers have certain level of education, and 43 percent have education up to secondary level. However, educational levels of husbands are lower than wives.

Around half of the mothers/caregivers watch television. Among those, more than 80 percent watch drama/serial and cinema. Radio is not popular. Only 3 percent mothers/caregivers in intervention and 9 percent in comparison areas use internet. Among the internet users, almost all use social media site, of which 50 percent use *Facebook*, followed by *Google*. Only 5 percent in intervention and 17 percent in comparison areas use SMS services in their mobile phones. And, only 9 percent household members have affiliation with microcredit organizations, such as- ASSA, Grameen Bank, and BRAC.

### Findings of the KAP survey with mothers/caregivers

**Diarrhoeal episode in under-five children:** Similar to the baseline, majority (95%) of under-five children in intervention and comparison areas have only a single episode of diarrhoea during last 30 days. However, the days of sufferings on an average are 3.6 days in intervention and 3.1 days in comparison areas at the end-line.

**Understanding on concept of diarrhea:** Concept of diarrhea (loose or liquid stools at least 3 times per day) is correctly known to 50 percent mothers/caregivers in intervention and 44 percent in comparison area, with net change 3.8 percentage points.

**Knowledge about ORS and its source:** All the mothers/caregivers are aware of ORS. However, around 30 percent still can't explain the method of ORS preparation. In public sector, Community Clinic (CC) is the primary source of ORS known to most of them in intervention (66%) and comparison (73%) areas. In private sector, *gram daktar* (village doctor) with pharmacy is the most popular source of ORS in intervention (75%) and in comparison (69%) areas during the end-line.

**Knowledge about zinc and its source:** Knowledge of the mothers/caregivers about zinc is fairly high at intervention (76%) than in comparison (44%) areas during end-line. It was lower at the baseline. The net change in this regard is 37.6 percentage points. And, the major known source of zinc known to them in public sector is CC (74.4% in intervention and 71.3% in comparison areas). In private sector, *gram daktar* with pharmacy is the major source of zinc known to them (69.3% in intervention and 35.6% in comparison areas).

**Knowledge about duration of treatment with zinc:** During the end-line, 40.4 percent mothers/caregivers in intervention and 23.4 percent in comparison areas reported correctly that, zinc is to be given for 10 days or more. The net change in this regard is 23.7 percentage points.

**Consultation for treatment of acute diarrhoea of under-5 children:** During the end-line, 35.3 percent mothers/caregivers in intervention sought treatment from public sector, and 61.3 percent from private sector for treatment of acute diarrhoea of their under-5 children. However, excluding gram daktar the private sector figure comes down to 23.2 percent, as 42.3 percent sought treatment from gram daktar.

**Prescription for treatment by service providers for acute childhood diarrhea:** Practice of advising to take both zinc and ORS with or without other medicine by service providers for treatment of acute diarrhea in children is 70.9 percent in intervention, and 37.0 percent in comparison areas, which were much lower at baseline.

**Coverage of zinc and ORS:** During end-line, 67.8 percent mothers/caregivers in intervention areas reports receipt of zinc and ORS from any source for treatment of acute diarrhea in children, of which 38.1 received from public sources and 55.2 from private sources. Coverage was much lower (27.6%) during baseline.

**Adherence to giving zinc tablet by caregivers for 10 days:** During end-line, 70.1 percent mothers/caregivers receiving zinc tablets from any sector reports giving it for 10 days for treatment of acute diarrhea in children in intervention areas. The same was 20.2 during baseline. Among those receiving zinc tablets from public sector 77.1 percent intervention areas reports giving it to their children for 10 days, which was only 21.1 percent during baseline.

**Adherence to giving zinc (tablet or syrup) by caregivers for 10 days:** During end-line, 53 percent mothers/caregivers receiving zinc from any sector reports giving it for 10 days for treatment of acute diarrhea in children in intervention areas. The same was 21 during baseline. Among those receiving zinc tablets from public sector 71.2 percent in intervention areas reports giving it to their children for 10 days, which was only 19.4 percent during baseline.

**Adherence to recommended dosage:** During end-line, 64.2 percent mothers/caregivers receiving zinc tablet and ORS from any sector reports giving zinc tablets for 10 days and 2 sachets of ORS for treatment of acute diarrhea in children in intervention areas. The same was 20.9 percent during baseline. Among those receiving zinc tablets and ORS from public sector 71.9 percent in intervention areas reports following this dosage, which was only 19.4 percent during baseline.

**Place of collection of ORS:** Private sector (75%) is still the most popular place for ORS collection to the mothers/caregivers as compared to public sector (39%), specially, *gram daktar* with pharmacy in intervention (52%) and in comparison areas (31%). In public sector, CCs are most popular (26%).

**Place of collection of zinc tablet and/or syrup:** Mothers/caregivers mostly collect zinc tablet from Community Clinic (64%), and syrup from *gram daktar* with pharmacy (61%) in intervention areas.

**Counseling:** According to mothers/caregivers, 66 percent of them in intervention and 33 percent in comparison areas they have been counseled by health workers to use zinc and ORS.

### **Findings of Interviews with Health Workers**

Two-thirds of the health workers are female. A high proportion of them (above 35%) are master degree holders. More than 50 percent of them received basic training more than 5 years back. Around 80 percent health workers in intervention and 9 percent in comparison areas report receipt of training on use of zinc for treatment of acute diarrhea in children from Integrated Management of Childhood Illness (IMCI) programme of DGHS, or MI. Around 79.8 percent health workers in intervention and 72.9 percent in comparison areas can spell out correctly that passage of at least three loose stools per day is diarrhea.

**Knowledge on zinc supplement:** The important advantages of giving zinc in acute diarrhoea in children are pronounced more than 80 percent service providers in intervention and a lower proportion in comparison areas. Around 91 percent health workers in intervention and 81 percent in comparison areas reports knowing the recommended dose of zinc for children aged 6-59 months. It was much lower during baseline. A 95 percent service providers in intervention and 70 percent in comparison areas knows correct duration of using zinc (10-14 days). Around 82 percent health workers in intervention and 61 percent in comparison areas were able to demonstrate the procedure of using dispersible zinc tablet correctly.

**Practices of prescribing zinc and ORS in management of childhood diarrhoea:** A 73 percent of health workers in intervention and 46 percent in comparison areas claims prescribing both zinc and ORS for management of childhood diarrhoea, which was lower at baseline.

**Adequate stock of zinc and ORS at present:** A 54 percent health workers in intervention and 49 percent in comparison areas report that they have adequate stock of ORS and zinc at present.

**Knowledge on estimating demand of zinc and ORS:** A 22 percent health workers in intervention and 31 percent in comparison areas report that they know how to estimate the demand of zinc and ORS in their area. However, 51 percent in intervention and 21 percent in comparison areas can demonstrate the procedure of calculation of it correctly.

**Household visit made by health workers:** Overall, 54 percent health workers during end-line reports that have claims visiting the houses of children during diarrhoea. However, the mothers/caregivers report that none of them visited their houses during that period.

**Conduction of group counseling session:** The health workers report that, 88 percent of them in intervention areas and 60 percent in comparison areas are conducting group counseling sessions with the mothers/caregivers. Overall, 98 percent health workers in intervention and 68 percent in comparison areas have discussed the benefit of zinc in group counseling session in the last month.

### **Findings of Interviews with Health Supervisors**

Health supervisors are dominated by females aged 45 years or more. A 94 percent health supervisor in intervention and 61 percent in comparison areas during end-line reports of their awareness about the use of zinc and ORS in the management of childhood diarrhea. Overall, 72 percent of them in intervention and 31 percent in comparison correctly know the dosage of 20 mg zinc tablet daily, and 94 percent in intervention and 46 percent in comparison areas know of correct duration of zinc tablet intake for 10-14 days. The health supervisors on an average report of undertaking at least one visit per month for supervision of use of zinc and

ORS in the management of diarrhea in under-5 children, around 84 percent of them use checklist during such supervision. Around 52 of supervisors in intervention and 42 percent in comparison areas report receiving complete monthly report/HMIS on use of zinc and ORS for diarrhea of under-5 children on time. In order to improve coverage and compliance of zinc and ORS, the health supervisors suggest having adequate supply of zinc tablets and ORS in all health facilities and health posts, and good counseling and clear information to the patients on benefit, dose, and administration of zinc and ORS.

### **Findings of Interviews with *gram daktars* (Village doctors)**

As high as 99 percent of the *gram daktars* in intervention and 88 percent in comparison areas perceived diarrhoea as at least three loose stools per day. However, 79 percent in intervention and 52 percent in comparison areas are aware of correct dose of zinc supplement (20 mg/day), and 84 percent in intervention and 41 percent in comparison areas are aware of correct duration of intake of zinc tablet for 10-14 days. Overall, 91 percent *gram daktars* in intervention and 50 percent in comparison areas can demonstrate preparation method of dispersible zinc tablet correctly. In general, 65 percent *gram daktars* in intervention and 36 percent in comparison areas have prescribed both zinc and ORS with or without other medicines. In order to improve coverage and compliance to use of zinc they suggest to establish a good relation with the mothers/caregivers, provide zinc tablets and ORS to all health facilities, provide adequate counseling to the mothers/caregivers on dosage, use, benefits and administration of zinc and ORS, and follow up home visits.

### **Findings of Interviews with Health Managers**

According to the health managers, total success of the programme is largely dependent on appropriate training of the frontline workers on zinc and ORS. According to them, the role of counseling by the frontline workers is very much effective. However they admitted that, none of the health managers is involved in monitoring of the MI initiated programme on scaling up the use of zinc and ORS in the treatment of diarrhea currently, and there is no special discussion on childhood diarrhea and challenges in providing zinc and ORS to every under 5 child with diarrhoea in the monthly coordination meetings held at UHCs. They also admitted that, they do not have any idea on IEC/BCC materials related to this programme. However, to achieve success in programme implementation they suggest- intensive training of health workers as well as refresher sessions regularly to provide both zinc and ORS to each under 5 children with diarrhoea; advocacy meeting with the other officials, local govt. representatives, community leaders, and religious leaders; effective and regular monitoring of the programme activities using checklist and other appropriate tools; ensuring adequate supply of zinc and ORS in every CCs, FWCs, and UHCs; establish regular home visit by the health workers and meeting with the mothers/caregivers; and inclusion of NGO health workers in this programme.